

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51904189>

Unmasking Masks in Makkah: Preventing Influenza at Hajj

Article in *Clinical Infectious Diseases* · January 2012

DOI: 10.1093/cid/cir826 · Source: PubMed

CITATIONS

16

READS

190

7 authors, including:



Harunor Rashid

Children's Hospital at Westmead

127 PUBLICATIONS 1,960 CITATIONS

[SEE PROFILE](#)



Robert Booy

The University of Sydney

465 PUBLICATIONS 14,101 CITATIONS

[SEE PROFILE](#)



Leon Heron

The New South Wales Department of Health

76 PUBLICATIONS 1,714 CITATIONS

[SEE PROFILE](#)



Ziad Memish

King Saud medical city

1,174 PUBLICATIONS 107,533 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



MERS-CoV outbreak control [View project](#)



Flu Watch [View project](#)

Unmasking Masks in Makkah: Preventing Influenza at Hajj

Each year more than 2 million people from all over the world attend the Hajj pilgrimage to Saudi Arabia. At least 60% of them develop respiratory symptoms there or during outward or homebound transit [1, 2]. During recent interpandemic years, approximately 1 in 10 pilgrims with respiratory symptoms in Makkah have had influenza detected by polymerase chain reaction tests of respiratory samples [3, 4]. Pneumonia is the leading cause of hospitalization at Hajj, accounting for approximately 20% of diagnoses on admission [5]. Hajj 2009 took place in November in the middle of the influenza A H1N1 pandemic (pH1N1). However, a recent report suggests that the

prevalence of pH1N1 was 0.2% ($n = 1$) among 519 arriving pilgrims and 0.1% ($n = 2$) among 2699 departing pilgrims [6], probably because of heightened infection control and preexisting cross-immunity among pilgrims, most of whom would have been exposed to H1N1 before 1957 [7].

Experience from an ear, nose, and throat (ENT) clinic in Makkah suggested that only 77 of 3087 patients (2.5%) who attended with ENT symptoms had suspected pH1N1 based on history of contact with a patient with pH1N1 and clinical symptoms of influenza-like illness (ILI); however, laboratory testing for influenza virus was not undertaken [8]. The efficacy of influenza vaccine is dependent on matching the vaccine and circulating influenza strains. Because the pH1N1 vaccine was not largely available in time for Hajj 2009, protection was dependent on previous exposure to H1N1 influenza A virus and nonpharmaceutical interventions such as face masks [2].

Historically, face masks have been used to prevent or reduce nosocomial transmission of pandemic influenza since, at least, the time of Spanish influenza in 1918. Nurses who wore specially designed face masks and changed them every 2 hours experienced lower infection rates than those who did not [9]. Recent experiments confirm that surgical masks and respirators can filter influenza virus, although observational studies or clinical trials have not yet clearly demonstrated the effectiveness of plain surgical masks in household or healthcare settings [10–16].

The role of surgical masks, respirators, and/or hand hygiene in preventing ILI has been examined in at least 15 studies including 5 randomized controlled trials (RCTs). These compared “plain surgical mask” with “no intervention” against ILI [10–14]. In 4 of these RCTs influenza was laboratory confirmed [10–13]. Metadata from these studies indicated that wearing surgical masks did not

change the frequency of laboratory-confirmed influenza (Figure 1). The findings may have been due to limitations and biases in study design [10–14]. A common limitation of all studies was small sample size. Despite similar study design, the sample-size calculations were based on different assumptions, and the studies lacked the power to detect a difference in incidence of laboratory-confirmed influenza.

At least 2 observational studies at the Hajj have examined the role of masks in preventing acute respiratory infection. One of these studies evaluated the role of face masks among pilgrims at the 2002 Hajj [15], when a protective effect was shown in men but not in women. The other study evaluated the use of face masks worn by healthcare workers at the 2005 Hajj in preventing acquisition of acute respiratory infection when protective effectiveness was non-significant [16].

The Hajj congregation of more than 2 million people who stay for 5 days in 25 000 tents in Mina, a valley within Makkah city, provides a unique opportunity to conduct RCTs of mask effectiveness to avoid sample size and design limitations.

During Hajj 2011 the 5-day stay in Mina is from 4 to 8 November. We are undertaking a pilot study to examine the feasibility of a cluster RCT of mask use among Australian pilgrims. Assuming success of this pilot trial, we plan to set up large RCT involving pilgrims from several countries in subsequent years. We are seeking expressions of interest for a multinational collaboration and would be pleased to hear from possible collaborators.

Note

Potential conflicts of interest. R. B. has received financial support from the pharmaceutical companies CSL, Sanofi, GlaxoSmithKline, Novartis, Roche, and Wyeth to conduct research and attend and present at scientific meetings. Any funding received is directed to a National Centre for Immunisation Research




Figure 1. Comparison between mask and control groups in randomized controlled trials. Abbreviations: CI, confidence interval; df, degrees of freedom; M-H, Mantel–Haenszel.

and Surveillance research account at The Children's Hospital at Westmead and is not personally accepted by R. B. L. H. conducts sponsored research for several vaccine manufacturers; has received support to attend conferences from GlaxoSmithKline and Sanofi Pasteur; and has been a member of a vaccine advisory board for Novartis. All funds received are directed to the National Centre for Immunisation Research and Surveillance of Vaccine Preventable Diseases, The Children's Hospital at Westmead. J. N.-V.-T. has received funding to attend influenza-related meetings, lecture and consultancy fees, and research funding from several influenza antiviral drug and vaccine manufacturers. All forms of personal remuneration ceased in September 2010, but influenza-related research funding from GlaxoSmithKline, F. Hoffmann-La Roche, and AstraZeneca remains current. He is a former employee of SmithKline Beecham plc (now GlaxoSmithKline), Roche Products Ltd, and Aventis-Pasteur MSD (now Sanofi-Pasteur MSD), all prior to 2005, with no outstanding pecuniary interests by way of shareholdings, share options, or accrued pension rights. All other authors report no potential conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

Harunor Rashid,¹ Robert Booy,^{1,2} Leon Heron,¹ Ziad A. Memish,³ Jonathan Nguyen-Van-Tam,⁴ Osamah Barasheed,¹ and Elizabeth Haworth⁵

¹National Centre for Immunisation Research and Surveillance, The Children's Hospital at Westmead, Sydney, and ²Sydney Emerging Infections and Biosecurity Institute, The University of Sydney, New South Wales, Australia; ³Ministry of Health, Riyadh, Saudi Arabia; ⁴Division of Epidemiology and Public Health, University of Nottingham (World Health Organization Collaborating Centre for Pandemic Influenza and Research), and ⁵The Academic Department of Public Health, Oxford University, United Kingdom

References

- Deris ZZ, Hasan H, Sulaiman SA, Wahab MS, Naing NN, Othman NH. The prevalence of acute respiratory symptoms and role of protective measures among Malaysian hajj pilgrims. *J Travel Med* **2010**; 17:82–8.
- Gautret P, Vu Hai V, Sani S, Douchi M, Parola P, Brouqui P. Protective measures against acute respiratory symptoms in French pilgrims participating in the Hajj of 2009. *J Travel Med* **2011**; 18:53–5.
- Rashid H, Shafi S, Haworth E, et al. Viral respiratory infections at the Hajj: comparison between UK and Saudi pilgrims. *Clin Microbiol Infect* **2008**; 14:569–74.
- Alborzi A, Aelami MH, Ziyaeyan M, et al. Viral etiology of acute respiratory infections among Iranian Hajj pilgrims, 2006. *J Travel Med* **2009**; 16:239–42.
- Madani TA, Ghabrah TM, Al-Hedaithy MA, et al. Causes of hospitalization of pilgrims in the Hajj season of the Islamic year 1423 (2003). *Ann Saudi Med* **2006**; 26:346–51.
- Memish ZA, Assiri AM, Hussain R, Alomar I, Stephens G. Detection of respiratory viruses among pilgrims in Saudi Arabia during the time of declared influenza A H1N1 pandemic. *J Travel Med* **2011** (in press).
- Xu R, Ekiert DC, Krause JC, Hai R, Crowe JE Jr, Wilson IA. Structural basis of preexisting immunity to the 2009 H1N1 pandemic influenza virus. *Science* **2010**; 328:357–60.
- Alherabi AZ. Impact of pH1N1 influenza A infections on the Otolaryngology, Head and Neck Clinic during Hajj, 2009. *Saudi Med J* **2011**; 32:933–8.
- Hobday RA, Cason JW. The open-air treatment of pandemic influenza. *Am J Public Health* **2009**; 99:S236–42.
- Aiello AE, Murray GF, Perez V, et al. Mask use, hand hygiene, and seasonal influenza-like illness among young adults: a randomized intervention trial. *J Infect Dis* **2010**; 201: 491–8.
- Cowling BJ, Fung RO, Cheng CK, et al. Preliminary findings of a randomized trial of non-pharmaceutical interventions to prevent

- influenza transmission in households. *PLoS One* **2008**; 3:e2101.
12. MacIntyre CR, Cauchemez S, Dwyer DE, et al. Face mask use and control of respiratory virus transmission in households. *Emerg Infect Dis* **2009**; 15:233–41.
 13. MacIntyre CR, Wang Q, Cauchemez S, et al. A cluster randomized clinical trial comparing fit-tested and non-fit-tested N95 respirators to medical masks to prevent respiratory virus infection in health care workers. *Influenza Other Respi Viruses* **2011**; 5:170–9.
 14. Canini L, Andreoletti L, Ferrari P, et al. Surgical mask to prevent influenza transmission in households: a cluster randomized trial. *PLoS One* **2010**; 5:e13998.
 15. Choudhry AJ, Al-Mudaimegh KS, Turkistani AM, Al-Hamdan NA. Hajj-associated acute respiratory infection among hajjis from Riyadh. *East Mediterr Health J* **2006**; 12: 300–9.
 16. Al-Asmary S, Al-Shehri AS, Abou-Zeid A, Abdel-Fattah M, Hifnawy T, El-Said T. Acute respiratory tract infections among Hajj medical mission personnel, Saudi Arabia. *Int J Infect Dis* **2007**; 11:268–72.

Correspondence: Harunor Rashid, MBBS, DCH, MD, National Centre for Immunisation Research and Surveillance, The Children's Hospital at Westmead, Cnr Hawkesbury Rd and Hainsworth St, Locked Bag 4001, Westmead, NSW 2145, Australia (harunorr@chw.edu.au).

Clinical Infectious Diseases 2012;54(1):151–3

© The Author 2011. Published by Oxford University Press on behalf of the Infectious Diseases Society of America. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

DOI: 10.1093/cid/cir826